

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SUPERWAY BROADLEAF BINDII & CLOVER SELECTIVE HERBICIDE

ACTIVE CONSTITUENTS: 280 g/L MCPA (present as the iso-octyl ester)
140g/L BROMOXYNIL (present as the octanoate ester)
40 g/L DICAMBA (present as the acid)
SOLVENT: 256 g/L HYDROCARBON LIQUID

GROUP	C I	HERBICIDE
--------------	------------	------------------

For the control of certain broadleaf weeds in turf and wheat, barley, commercial rye, oats, triticale as per the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS 1L, 2.5L, 5L, 10L, 20L, 200L, 1000L

SUPERWAY GARDEN PRODUCTS PTY LTD
Shed 6, Cnr Roys Road & Moroney Place
BEERWAH QLD 4159
ABN 24 056 553 274
TEL (07) 54390355

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact or inhaled or swallowed. Will irritate the eyes and skin. Do not inhale vapour or spray mist. Repeated exposure may cause allergic disorders. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PVC gloves, face shield and goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow length PVC gloves. If product in eyes, wash out immediately with soap and water.

After each day's use wash gloves, face shield and goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 131126).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

CONDITIONS OF SALE

The use of Superway Broadleaf, Bindii & Clover Selective Herbicide is beyond the control of the Superway Garden Products Pty Ltd and no warranty, expressed or implied, is given by Superway Garden Products Pty Ltd regarding its suitability or efficacy for any purpose for which it is used by the buyer, except the non-excludable rights under State/Territory or Federal legislation.

APVMA Approval No: 58780/1L/0504

58780/2.5L/0504

58780/5L/0504

58780/10L/0504

58780/20L/0504

58780/200L/0504

58780/1000L/0504

DOM:

Batch No:

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

SUPERWAY BROADLEAF BINDII & CLOVER SELECTIVE HERBICIDE

**ACTIVE CONSTITUENTS: 280 g/L MCPA (present as the iso-octyl ester)
140g/L BROMOXYNIL (present as the octanoate ester)
40 g/L DICAMBA (present as the acid)
SOLVENT: 256 g/L HYDROCARBON LIQUID**

GROUP	C I	HERBICIDE
--------------	------------	------------------

For the control of certain broadleaf weeds in turf and wheat, barley, commercial rye, oats, triticale as per the Directions for Use table.

THIS LEAFLET IS PART OF THE LABEL

APVMA APPROVAL NO: 58780/0504

**SUPERWAY GARDEN PRODUCTS PTY LTD
Shed 6, Cnr Roys Road & Moroney Place
BEERWAH QLD 4159
ABN 24 056 553 274
TEL (07) 54390355**

DIRECTIONS FOR USE:

RESTRAINTS: DO NOT spray when rain seems likely to occur within 3 hours.

DO NOT spray when weeds are wet with dew or rain, or under stress from drought, low soil fertility, extreme cold or water-logging. DO NOT spray when crop is under stress from drought, excessively wet conditions or low soil fertility.

DO NOT apply to crops undersown with legumes.

CROP	CROP STAGE	WEEDS	STATE	RATE	CRITICAL COMMENTS
Recreational Turf, Established Couch (including Queensland Blue), Kikuyu, Paspalum, Fescue, Bent and Rye Lawns		Broadleaf weeds including: Bindy-Eye (<i>Calotis hispidula</i>), Capeweed (<i>Arctotheca calendula</i>), Carrot Weed (<i>Cotula australis</i>), Catsear (<i>Hypochoeris radicata</i>), Caustic weed (<i>Euphorbia drummondii</i>), Clovers (<i>Trifolium</i> spp.), Common Plantain (<i>Plantago</i> spp.), Creeping buttercup (<i>Ranunculus repens</i>), Cudweed (<i>Gnaphalium</i> spp.), Dandelion (<i>Taraxacum officinale</i>), Fleabane (<i>Ictonyza</i> spp.), Funnel Weed (<i>Cotula turbinata</i>), Jo-Jo (Onehunga) (<i>Salvia pterisperma</i>), Thistles, Three cornered Jack (Doublegee) (<i>Emex australis</i>), Wood Sorrel (<i>Oxalis</i> spp.)	NSW, Qld, SA, Vic, WA, ACT only	4-5L in 150L of water per hectare	Apply to actively growing weeds. A slight browning of turf, especially Paspalum, may occur under hot conditions. Use the lower rate for weeds up to the 4 leaf stage. DO NOT mow turf for 2 days after treatment. DO NOT apply to Buffalo Grass. DO NOT feed turf/lawn clippings from treated areas to poultry or livestock.
Wheat, Barley, Oats, Triticale, Commercial Rye	3 leaf to fully tillered from Zadocks (13 to 30)	Group A**: Ball Mustard (<i>Neslia paniculata</i>), Capeweed (<i>Arctotheca calendula</i>), Docks, seedlings only (<i>Rumex</i> sp.), Doudle gee (Spiny emex, Three cornered Jack) (<i>Emex australis</i>), Hedge Mustard (<i>Sisymbrium officinale</i>), Indian Hedge Mustard (<i>Sisymbrium orientale</i>), Wild Mustard (<i>Sisymbrium</i> sp), Wild Radish (<i>Raphanus raphanistrum</i>), Wild Turnip (<i>Brassica tournefortii</i>), Wireweed (Hogweed) <i>Polygonum aviculare</i> , Yellow burr Weed (<i>Amsinkia</i> (<i>Amsinkia</i> sp)	NSW, ACT, Vic, SA, WA only	750mL/ha	3 to 4 leaf stage but no more than 3cm in diameter.
				1L/ha	4 to 6 leaf stage but not more than 5cm in diameter.
	5 leaf to fully tillered (from Zadocks 15 to 30)	Turnip Weed (<i>Rapistrum rugosum</i>), Rough Poppy (<i>Papaver hybridum</i>), Saffron Thistle (<i>Carthamus lanatus</i>) Deadnettle (<i>Lamium amplexicaule</i>), Vetch (<i>Vicia</i> spp.) Charlock (<i>Sinapsis arvensis</i>), Fat Hen (<i>Chenopodium album</i>), Lesser Swinecress (Bitter cress) (<i>Coronopus didymus</i>)	WA only	1.4L/ha	6 to 8 leaf stage but not more than 10cm in diameter.
			Vic only		
			Tas only		
3 leaf to fully tillered (from Zadocks 15 to 30cm)	Group B** Climbing Buckwheat (Black bindweed)(<i>Polygonum convolvulus</i>), Common Iceplant (<i>Mesembryanthemum crystallinum</i>), Dense Flowered Fumitory – Red (<i>Fumaria densiflora</i>), Hoary Cress (<i>Cardaria draba</i>), Small Flowered Fumitory – White (<i>Fumaria parviflora</i>), Tree Hogweed (<i>Polygonum patulum</i>), White Iron weed, (Sheepweed, Corn Gromwell (<i>Buglossoides arvensis</i>) Slender Thistle (<i>Carduus</i> spp.), Spear Thistle (<i>Cirsium vulgare</i>), Variegated Thistle (<i>Silybum marianum</i>)Wall Fumitory (Pink weed) (<i>Fumaria muralis</i>)	NSW, ACT, Vic, SA, WA only	1L/ha	2 to 4 leaf stage but no more than 3cm in diameter.	
		1.4L/ha	6 to 8 leaf stage but not more than 10cm in diameter		
5 leaf to fully tillered (from Zadocks 15 to 30)		Tas only	1.4L/ha	4 to 6 leaf stage but not more than 100-120mm in diameter. Wall Fumitory is severely stunted by 1.4L/ha.	
**Where GROUP A & B weeds occur together, use the highest application rate.					

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: DO CUT OR GRAZE FOR STOCKFOOD FOR 14 DAYS AFTER APPLICATION.

General Instructions

Note: A temporary wilting may be evident in crops after application. Under some conditions, regrowth and successive germination of weeds may occur following treatment and may require follow-up spraying.

Crop Safety:

DO NOT spray outside crop growth stages recommended as crop damage may result. DO NOT apply to recently released cereal varieties until advice has been sought from the manufacturer or Local Department of Agriculture.

Mixing:

Add the required amount of the product directly to the water in tank and mix well.

Application:

DO NOT apply by misting machines.

For Boom application: Apply by a properly calibrated boom spray using not less than 50 litres of water per hectare.

For Aerial Application: Apply at least 25 litres of total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.

RESISTANT WEEDS WARNING

GROUP	C I	HERBICIDE
--------------	------------	------------------

Superway Broadleaf, Bindii & Clover Selective Herbicide is a member of the nitriles, phenoxy and benzoic acids groups of herbicides. Superway Broadleaf, Bindii & Clover Selective Herbicide has the inhibitors of photosynthesis at photosystem II and disruptors of plant cell growth mode of action. For weed resistance management, Superway Broadleaf, Bindii & Clover Selective Herbicide is a Group C and Group I herbicide. Some naturally occurring weed biotypes resistant to Superway Broadleaf, Bindii & Clover Selective Herbicide and other Group C and Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Superway Broadleaf, Bindii & Clover Selective Herbicide or other Group C and Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Superway Garden Products Pty Ltd accepts no liability for any losses that may result from the failure of Superway Broadleaf, Bindii & Clover Selective Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Spray equipment must be calibrated accurately prior to working on crops. Keep container sealed (airtight) when not in use. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops.

DO NOT use in close proximity to vines, tomatoes and other susceptible crops including flowers, vegetables, ornamentals, lupins and fruit trees. To prevent damage, DO NOT permit drift or vapour or spray to come into contact with fruit and ornamental trees, cotton, vines, vegetables, flowers, tobacco, lupins or any other field crop. Minimise spray drift by using low pressure and nozzles which do not give a fine droplet size.

DO NOT spray in high winds. DO NOT use this product on or near desirable plants or trees or in locations where the chemical may be washed into contact with their roots.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT use container for any other purpose. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact or inhaled or swallowed. Will irritate the eyes and skin. Do not inhale vapour or spray mist. Repeated exposure may cause allergic disorders. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PVC gloves, face shield and goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow length PVC gloves. If product in eyes, wash out immediately with soap and water.

After each day's use wash gloves, face shield and goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 131126).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

CONDITIONS OF SALE

The use of Superway Broadleaf, Bindii & Clover Selective Herbicide is beyond the control of the Superway Garden Products Pty Ltd and no warranty, expressed or implied, is given by Superway Garden Products Pty Ltd regarding its suitability or efficacy for any purpose for which it is used by the buyer, except the non-excludable rights under State/Territory or Federal legislation.